

LETTER • OPEN ACCESS

Human dependence on natural resources in rapidly urbanising South African regions

To cite this article: Stefano Balbi *et al* 2019 *Environ. Res. Lett.* **14** 044008

View the [article online](#) for updates and enhancements.

You may also like

- [Investigation of the speed regime of tractor diesel engine running on natural gas with recirculation](#)
V A Likhanov and O P Lopatin
- [Spectral representations of topological groups and near-openly generated groups](#)
V. M. Valov and K. L. Kozlov
- [Study on dynamic mechanical properties of magnetorheological elastomers based on natural rubber/thermoplastic elastomer hybrid matrix](#)
Xincheng Song, Wenju Wang, Fufeng Yang *et al.*

Recent citations

- [The journey to monitoring ecosystem services: Are we there yet?](#)
Ana Sofia Vaz *et al*
- [Environmental Services: A New Approach Toward Addressing Sustainable Development Goals in Sub-Saharan Africa](#)
Jasper Knight
- [Tracing and building up environmental justice considerations in the urban ecosystem service literature: A systematic review](#)
Amalia Calderón-Argelich *et al*



IOP Publishing

ENVIRONMENTAL RESEARCH 2021

A VIRTUAL CONFERENCE
15–19 NOVEMBER

FREE TO
ATTEND

REGISTER
NOW

Environmental Research Letters



LETTER

OPEN ACCESS

RECEIVED
19 July 2018

REVISED
20 December 2018

ACCEPTED FOR PUBLICATION
14 January 2019

PUBLISHED
29 March 2019

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Human dependence on natural resources in rapidly urbanising South African regions

Stefano Balbi^{1,8} , Odirilwe Selomane^{2,3,8,9} , Nadia Sitas^{3,4}, Ryan Blanchard^{3,5}, Ilse Kotzee³, Patrick O'Farrell^{3,6} and Ferdinando Villa^{1,7}

¹ Basque Centre for Climate Change (BC3), Sede Building, Campus EHU/UPV, Leioa, Bizkaia, Spain

² Stockholm Resilience Centre, Stockholm University, Sweden

³ Council for Scientific and Industrial Research, South Africa

⁴ Department of Conservation Ecology and Entomology, Stellenbosch University, South Africa

⁵ Centre for Invasion Biology, Stellenbosch University, South Africa

⁶ Percy FitzPatrick Institute of African Ornithology, University of Cape Town, Rondebosch, South Africa

⁷ IKERBASQUE, Basque Foundation for Science, Bilbao, Bizkaia, Spain

⁸ Joint first authors.

⁹ Author to whom any correspondence should be addressed.

E-mail: onebrownbread@gmail.com

Keywords: provisioning ecosystem services, sustainable development, urban transition, machine learning, openly accessible data, informality


Abstract

Enhancing the governance of social-ecological systems for more equitable and sustainable development is hindered by inadequate knowledge about how different social groups and communities rely on natural resources. We used openly accessible national survey data to develop a metric of overall dependence on natural resources. These data contain information about households' sources of water, energy, building materials and food. We used these data in combination with Bayesian learning to model observed patterns of dependence using demographic variables that included: gender of household head, household size, income, house ownership, formality status of settlement, population density, and in-migration rate to the area. We show that a small number of factors—in particular population density and informality of settlements—can explain a significant amount of the observed variation with regards to the use of natural resources. Subsequently, we test the validity of these predictions using alternative, open access data in the eThekweni and Cape Town metropolitan areas of South Africa. We discuss the advantages of using a selection of predictors which could be supplied through remotely sensed and open access data, in terms of opportunities and challenges to produce meaningful results in data-poor areas. With data availability being a common limiting factor in modelling and monitoring exercises, access to inexpensive, up-to-date and free to use data can significantly improve how we monitor progress towards sustainability targets. A small selection of openly accessible demographic variables can predict household's dependence on local natural resources.

1. Introduction

As countries develop and many societies transition to urbanisation, they tend to reduce their dependence on the local natural environment to meet basic needs (Anderson 1987, Cumming *et al* 2014, Sanderson *et al* 2018). This decoupling is often described by the transition from a strong reliance on agriculture as the main source of national income to other, less directly coupled sectors such as industrial and service sectors

(Daunton 1995, Mellor 1995, Soubbotina and Sheram 2000). Beyond agriculture, this transition entails a move from 'green-loops', i.e. social-ecological systems characterized by high dependence on local provisioning ecosystem services (ES), to 'red loops', characterised by intensive imports or mediated (see figure 1) provisioning ES from local or other systems (Cumming *et al* 2014). A key driver of this transition is urbanisation (Grimm *et al* 2018). Urban areas will continue to experience rapid increases of population, consumption,



and diet changes, among other factors into the future. How cities balance rapid urbanisation in relation to resource consumption may determine whether or not many of the sustainable development goals (SDGs) are achieved (UN 2015).

In today's globalized world, most of the urban consumption of natural resources is mediated by biophysical features (e.g. cross border river systems), formal and informal markets, international governance mechanisms (e.g. protection of threatened species) and infrastructures (e.g. dams) (Syrbe and Walz 2012, Reyers *et al* 2013, Palomo *et al* 2016). As a result, most people in urban environments consume natural resources produced or extracted somewhere else (Liu *et al* 2013, Nel *et al* 2017). However, cities are not homogenous entities and seeing them as such can mask their unique development trajectories and the differentiated dependences of various groups and sectors on local ecosystems. For example, urbanisation in the Global South is partly characterised by informality both in land-use, resource access, governance processes (Brouwer *et al* 2009, McFarlane 2012, Güneralp *et al* 2018). In-migration from rural areas usually occurs more quickly than infrastructure can be developed to meet the growing demand (Foster and Briceno-Garmendia 2010, Brueckner and Lall 2015). Additionally, there is incomplete and unequal access to the main infrastructure grid and other formally

provided services such as water and energy, which often imply that people have to rely on local ES as alternatives.

Urbanisation in cities in the Global South also involves a combination of people relocating permanently to cities, people moving to cities but maintaining strong connections to rural areas, and 'circular migration' which refers to repeated temporary migration between two or more areas (Collinson *et al* 2007, UNDP and Newland 2009, Masterson *et al* 2017). Urban dwellers in these cities continue to rely on local natural resources to satisfy basic needs (McHale *et al* 2013). This means that natural resources within the city boundaries play a broader role beyond providing recreational and cultural services, which are often the most commonly cited and studied urban ES (Bertram and Rehdanz 2015, La Rosa *et al* 2016). The direct extraction of natural resources within the city boundaries has implications for how natural resources in urban areas are managed and sustained.

Tools and approaches that can account for multiple modalities of reliance on the environment can help understand and manage natural resources and man-made infrastructure and support targeting of policies towards the SDGs. In contrast to the common framing of natural resource dependence as a dichotomous category, we propose a conceptualization of dependence as a gradient (figure 1). This builds on Cumming

et al (2014) and Hamann *et al* (2015)'s conceptualisations of 'green and red loops' which shows dependence on natural resources being both direct and indirect, or local and nonlocal. Similar to Raudsepp-Hearne *et al* (2010) and Hamann *et al* (2015), we use national survey data, which they used successfully to explore dependence on and use of ES at national and subnational scales. First, we develop a metric of households' overall dependence on natural resources based on Statistics South Africa's census data. Second we use households' demographic characteristics to model the same patterns observed in the census data using machine learning. Lastly, we explore whether similar patterns can be observed using other publicly available data. Our context of analysis is the eThekweni Metropolitan Municipality, comprising the city of eThekweni (Durban) in South Africa. We show that our simulation accurately models the empirical data and that some of the main variables driving the model can be sourced from alternative data sources in the absence of national statistics. Lastly, we discuss the implication of these findings for other data-scarce African metropolitan areas, and more generally for similar contexts in the Global South.

2. Materials and methods

2.1. Study area

The eThekweni Municipality is located on the east coast of South Africa in the province of KwaZulu Natal (figure 2). It spans an area of 2297 km² characterised by coastal plains and a steep, dissected topography. The municipality is a middle-income metropolitan region of 3.6 million people residing under highly unequal social, economic and environmental conditions. The urban landscape consists of commercial buildings, formal and informal settlements and peri-urban agricultural land. Approximately 68% is considered rural, with pockets of dense settlements, and about 25% of the inhabitant live in informal and rural settlements. The region therefore shares an urban and rural landscape with a wide range of settlement types, where both rural and urban areas experience significant population growth. The unit of analysis for the study was the municipal ward level, the smallest administrative unit in South Africa; there were 103 wards in the 2011 census, and 113 in 2017 due to changes in administrative borders. Wards are also the scale at which socio-demographic data are collected in the national census of South Africa (StatsSA 2011).

2.2. Model design

Our approach is articulated in four steps. First, we take advantage of the existing and openly accessible 2011 Census data from Statistics South Africa (<http://statssa.gov.za/>) to develop a metric of overall *direct* dependence on natural resources (equation (1)). This

metric of ES dependence, is determined by the product of the proportions of households in a ward that do not directly depend on the 5 natural resources considered (table 1). In this formulation, 100% means that all households in a particular ward depend on at least one provisioning ES, and 0% means that none of the households depend on any of the provisioning ES. The five ES were chosen for two main reasons: first, because they are provided by the census data, and second because they can be disaggregated to the smallest administrative unit in South Africa. Other data, e.g. water quality determined by households treating water before using it, are only available at the provincial scale, which is not a useful scale for our analysis.

$$ES\ Dep. = 100 * (1 - ((1 - pW) \cdot (1 - pCR) \cdot (1 - pH) \cdot (1 - pCF) \cdot (1 - pB))). \quad (1)$$

Secondly, we use seven socio-demographic variables available at the ward level in the South African Census data and one variable representing the coverage of open spaces across the municipality (described in table 2), to explain the output from the metric. Some of these variables were also used in previous literature (Hamann *et al* 2015). The socio-demographic factors we used in this study have been identified elsewhere in literature to be important determinants of how communities, households or individuals interact with the natural environment and ultimately how they benefit from it (Leach *et al* 1999, Daw *et al* 2011, Al-assaf *et al* 2014, Tadesse *et al* 2014, Yang *et al* 2015). The variable E6, which denotes the availability of natural open spaces within an administrative unit, was chosen as an environmental variable to determine whether or not open spaces matter for natural resources dependence.

We investigate the explanatory power of the input variables (E1–E8) by applying machine learning (Willcock *et al* 2018). Our learning model uses the WEKA (Waikato Environment for Knowledge Analysis) machine learning software (<http://cs.waikato.ac.nz/ml/weka/>) and integrated into k.LAB, the software powering the ARIES (Artificial Intelligence for Ecosystem Services; <http://aries.integratedmodelling.org/>) technology adopted for the study. We apply a Bayesian classifier (*BayesNet*) that learns a *naïve* Bayesian network from the data using the K2 algorithm with all input nodes connected to the output node (Witten *et al* 2018). The training process involved 10 cross-validation iterations, each using 90% of the data to train the model and 10% to validate it. The machine learning algorithm used quantitative variables, discretised in 10 equal intervals, for both inputs and outputs.

To estimate dependence in the future, we use projected population growth data in the 25 years between 2011–2036 as input for the trained model. Population growth was based on the annual growth rate of population during 2001–2011, which averaged 1.09%,





Table 1. Provisioning ES used to develop the metric. Dependence is determined as proportion of households using these natural resources as their main sources.

Provisioning ES	Natural resource	Key (equation (1))
Freshwater use	Water from rivers and springs	pW
Crop production	Crop production in communal areas	pCR
Solid fuels for heating	Firewood and charcoal	pH
Solid fuels for cooking	Firewood and charcoal	pCF
Natural materials for constructing houses	Wood, grass, soil	pB

Table 2. Explanatory variables used to model dependence on ecosystem services.

Variable	Description	Mean (SD)
(E1) Proportion of household with female head	Households headed by women	40.64 (6.49)
(E2) Proportion of large households	Households with more than 4 members	25.51 (10.65)
(E3) Proportion of poor households	2011 poverty line was ^a 620 ZAR/person/month (StatSA 2012)	59.32 (17.22)
(E4) Proportion of households owning their house of residence	Census data distinguishes between owning, renting and free occupation.	55.70 (6.49)
(E5) Proportion of households living in informal settlements	Unplanned settlements where housing is not in compliance with current planning and building regulations	14.82 (14.37)
(E6) Proportion of land covered by the Metropolitan Open Space Network (D'MOSS)	D'MOSS is a system of open spaces (land and water), that incorporates areas of high biodiversity value which are linked together in a viable network.	21.55 (15.19)
(E7) Population density	Hectares of land per household	2814.89 (5244.25)
(E8) Proportion of in-migration	People moving into the wards from other municipalities and beyond.	14.72 (10.91)

^a 1 USD was 8.1285 ZAR in 2011.



resulting in population growth from 3.4 million people in 2011 to 4.5 million people in 2036. The assumption of a constant growth rate is an obvious simplification given the dynamic migration, birth rates and other factors affecting the population. However, such approximate pictures of future dependence can still be informative and are commonly employed in ‘business as usual’ scenarios (Millennium Ecosystem Assessment 2005, Hunt *et al* 2012).

As a last step, we reapply the model at the pixel level, rather than at the ward level, using only two of the eight exploratory variables (table 2: E5, E7), simulating a scenario in which only global and remotely sensed data are available. We discuss the predictions of this ‘data-poor’ model scenario both in the eThekweni Municipality and in the Cape Town metropolitan region.

3. Results

3.1. Overall dependence metric

Figure 4 shows our metric of dependence, the percentage of households dependent on at least one of the five ES described in table 1, at the ward level. Around eThekweni, only a small proportion of households are directly dependent on one or more of the above listed natural resources, as one would expect of an urban setting. Such proportions increase the farther away from the city, as would also be expected. In the most marginal areas, 100% of households are directly accessing at least one natural resource among water, food (own grown), solid fuel (for heating and for cooking) and building materials.

3.2. Modelling dependence with socio-demographic variables

The machine learned Bayesian model, using eight input variables in table 2, is able to reproduce the empirical output with significant accuracy as reported in table 3: 92% of the instances are correctly classified using the training dataset and 60% using only the validation dataset. The spatial results are illustrated in figure 5(A). However, not all of the variables contribute equally. For example, a similar network using

only the predictors that describe population (E2, E5, E7, E8) is able to capture 80% of correct instances (using the training set). The influence analysis of the network shows that three out of eight predictors—population density, proportion of large households, and proportion of households living in informal settlements—are driving the results (see table 4). The strength of influence is calculated from the conditional probability tables and expresses the difference between the probability distributions of two nodes by looking at the posterior probability distribution of a node, for each possible state of the parent or child node. To summarize this difference, we report normalized Euclidean distance, although other types of distances (e.g. Hellinger) are also used (Anderson 2006).

The 25-year projected analysis shown in figure 5(B) suggests a slow and mixed urbanisation transition in the metropolitan region, with most wards only slightly reducing their dependence on local natural resources and few moving in the opposite direction. The first outcome is more evident in the coast and in the closest outskirts of the city, while the second is seen in peri-urban areas and in the interior, confirming a divide between more and less urbanised areas. This can be attributed to densification in formal areas.

3.3. Application of the model in data-scarce situations

As a preliminary test, we applied the machine learned model described previously using only two out of eight input variables: presence of informal settlements (E5) and population density (E7). These variables, producing the strongest influence on the model results, can easily be provided through alternative data sources, including remote sensing and open data (e.g. for informal settlements see Busgeeth *et al* 2008, Chakraborty *et al* 2015). The Bayesian model was run with vector files containing polygons for the informal settlements (provided by the regional authority for eThekweni and downloaded from an open data project on informal settlements, <http://ismaps.org.za>, for Cape Town). Population density data came from the global grid dataset provided by NASA (Gridded Population of the World: GPW, v4), referring to the year 2015. The model




Figure 4. Percentage of households directly dependent on at least one provisioning ES according to 2011 Census data (k.LAB screenshot).

Table 3. Machine learning cross-validation results on training set.

Evaluation metrics ^a	Using training set	10 folds cross-validation	Only validation set
Correctly classified instances ^b	95	70	6
Incorrectly classified instances	8	33	4
Kappa statistic	0.8458	0.2737	0.3939
Mean absolute error	0.0328	0.0771	0.0916
Root mean squared error	0.1134	0.2214	0.2589
Relative absolute error	29.5152%	68.7247%	75.5794%
Root relative squared error	49.4312%	96.1668%	103.2831%

^a All input nodes and the output node of the network are discretised in 10 equal intervals.

^b There were four incorrectly classified instances—a 20% class was classified as 10%, a 50% class as 40%, a 100% class as 80% and finally a 60% class as 100%.

was run in two spatial contexts, eThekweni and Cape Town metropolitan regions (figure 6), at 100 m resolution, although population density is currently offered at a resolution of 30 arc-seconds (approximately 1 km at the equator). The latter is thus the main limiting factor in achieving finer-scale results. In each 100 m cell, the Bayesian network model simulates expected output using data-based evidence for the two mentioned variables and the prior probabilities learned from the previous steps for the remaining variables.

Figure 6 highlights how the distinction among wards is lost in favour of a more evenly distributed outcome. The gradient of dependence is still visible,

particularly for the eThekweni context where the data about informal settlements is more comprehensive. Differently from eThekweni, the metropolitan area of Cape Town is regarded at 99% urban area.

4. Discussion

Given our results, we hypothesize that our data-driven model could be applied in other areas of South Africa or even in parts of the African continent, with fewer input variables, and still retain the ability to capture the dependence of households in metropolitan areas




Figure 5. Machine learned percentage of households directly dependent on at least one provisioning ES before (a) and after (b) a scenario of population growth simulating 25 years (images exported from k.LAB automatically generated report).

Table 4. Strength of influence by node measured with normalised Euclidian distance. The first four variables are the most important factors.

Variable	Average	Maximum
(E7) Population density	0.243 381	0.642 193
(E5) Proportion of households living in informal settlements	0.204 627	0.380 318
(E8) In-migration as a proportion of households in a ward	0.172 172	0.353 102
(E2) Proportion of large households	0.211 939	0.342 179
(E3) Proportion of poor households	0.177 729	0.318 287
(E6) Proportion of land covered by D'MOSS	0.183 719	0.296 675
(E4) Proportion of households owning their house of residence	0.182 403	0.287 565
(E1) Proportion of household with female head	0.176 141	0.268 139

on natural resources. However, although the model provides reasonable indications in the eThekweni and Cape Town cases, its exportability to other cases requires further investigation. At the same time, the combination of machine learning techniques and heterogeneous typologies of data can be customized to local needs and provide useful indications in data-scarce situations. Nevertheless, we showed that a simple model using 2 variables, is able to reasonably capture the spatial distribution of the gradient's extremes (respectively urban areas and natural areas)

and some key hotspots of transition coinciding with the presence of informal settlements. It is important to note that the conditions for how households in a particular geographic location depend on direct natural resources is also dependent on the availability, climatic conditions and other contextual factors.

While South Africa provides detailed demographic data at the ward level, this is not the norm in other parts of the continent (Chandy 2013, Randall and Coast 2015). A methodology capable of substituting or complementing incomplete or missing national survey data with freely available or less expensive information can improve the coverage of studies of natural resource dependence—a valuable outcome as most regions of the world lack such data (Karp *et al* 2015). Availability of such data is increasingly important for the governance of social-ecological systems, as countries and regions look for ways to achieve equitable and sustainable development.

Our study exemplifies the use and integration of multiple data sources to respond to multidimensional problems (Tallis *et al* 2012). More generally, this work points at a possible future where the role of census data changes from being the main source of information to supporting learning and validation of models that can be run with proxy inputs. The use of census data as a complement to other data has long been done for Small Area Estimation (Rao and Yu 1994, Pfeffermann 2002). Other combinations of data include use of




Figure 6. Application of the machine learned model with only two spatially explicit input variables in Cape Town (on the left) and eThekweni (on the right) metropolitan regions (k.LAB screenshots). (The scale ranges from low dependence (blue) to high dependence (red)).

night light coverages (Xie *et al* 2015) to estimate poverty in Africa, where direct poverty data are notoriously lacking (Serajuddin *et al* 2015, Selomane *et al* 2015). With the increased availability of remote sensing data (De Araujo Barbosa *et al* 2015, Cord *et al* 2015, 2017), it is becoming increasingly possible to assess changes over time at significantly higher frequencies. Integration technologies such as k.LAB (<http://integratedmodelling.org>), can reduce data preparation time and enable large-scale replication without additional costs by enabling rapid data and model access. Such advantages increase when the number of explanatory variables is small, as our study highlighted. With rapid assessment tools and easy web access to data and models, it becomes possible to envision monitoring services such as early warning systems, where specific structural transitions can be predicted based on continuously updated, remotely sensed data. In addition, this methodology could also be used by national statistic departments to track progress towards SDGs.

Our model has four main demographic driving variables (E2, E5, E7, E8) which overshadow the role of income. We expected income to be important, after studies (Shackleton *et al* 2008, Yang *et al* 2013) that highlight poverty as a significant determinant of dependence on natural resources. One explanation for this discrepancy is that South African households are shielded from much of the direct dependence because the government provides some free basic services to low-income households who cannot afford them,

such as water, sanitation, waste removal and electricity (Muller 2008, Bhorat *et al* 2012). Another explanation might involve the intricate connections between poverty and other variables (Unterhalter 2012, Djoudi *et al* 2016) such as gender, geographic positionality, ethnicity, immigration status etc. Additionally, poverty is also linked to and influenced by external drivers, path dependencies and cross-scale interactions (Haider *et al* 2018). As a result, its effects may be masked or embedded in other process, especially those associated with power and with how systems of oppression and marginalisation play out in informal settlements (Rao *et al* 2017).

Our analysis shows how investigating dependence with a fine-grained approach (e.g. ward level versus fine-resolution grids) unveils fine-grained variations not captured by census data aggregated by administrative unit. For example, both the eThekweni and the Cape Town metropolitan regions were previously classified as uniformly ‘red loop’, representing low direct dependence on local natural resources (Hamann *et al* 2015); our analysis shows variations in dependence, comprising low, high and transition loops. Within these areas we are now able to identify potential *transition hot-spots*, crucial to understanding key social-ecological dynamics, such as inequality of access to natural resource and public infrastructure. Such areas are shared by people relying on direct harvest of natural resources, municipal services, or on a combination of both, with a mixture of *green* and *red loops* across households. This suggests that dependence on natural

resources cannot be characterised in discrete groups, but rather along a gradient with underlying feedbacks as to when and how long natural resources are used. While our model is unable to determine which resources are preferred in space or time which still warrants a deeper analysis, identifying this gradient improves our ability to spot source areas of direct resource use. We argue that spatially explicit analysis based on verified criteria and finer-grained spatial data is an improvement over the use of data aggregated per administrative unit.

The lack of finer-scale temporal analysis is certainly an area in need of improvement (Karp *et al* 2015). Our attempt to deal with this limitation through projection of population growth rates does not suggest radical changes in natural resource dependence, which remains largely mixed (figure 5). It is possible that the level of dependence will persist for the foreseeable future (McHale *et al* 2013), as the importance of informality in the urban and peri-urban system is becoming a common characteristic of rapidly growing cities on the African continent, and generally in the Global South (Roy 2009, McFarlane 2012). Thus a key priority for the local authorities should be to maintain adequate local ecosystem health (e.g. by further preserving and/or expanding conservation areas like the D'MOSS) around the identified transition hotspots, so that the supply of ES can be maintained over time. We regard this as a robust policy recommendation even considering the assumption of constant population growth over time, which might be underplaying more complex dynamics to come. For example, dependence on local ecosystems is determined by more than population growth and may be mediated or substituted by municipal provision of services. Additionally, the urbanisation process is not linear and may even yield surprising and potentially positive outcomes (Sanderson *et al* 2018), as people move between cities and rural areas in a circular migration pattern (Collinson *et al* 2007, Mchale *et al* 2013). Yet this finding does highlight the need for adequate holistic management processes that are mindful of both human and environmental needs, and suggests that natural resources within these systems need to be managed accordingly along the gradient.

5. Conclusion

Survey-driven social data such as those made available by national statistical offices are expensive to gather and update frequencies are low (usually 10-year cycles). For example, much of the demographic data required to monitor key proxy indicators, such as poverty, remain difficult to access or too expensive to develop for many areas of the globe. Such information is likely to be patchy or inconsistent at a certain spatial scale, or available for very few points in time over long intervals. Thus, efforts to complement, harmonize

and validate inconsistent information and improve spatial and temporal resolution of such data products by means of more easily obtained data are of paramount importance. In this study we show that machine learning techniques can contribute to filling knowledge gaps concerning the use of natural resources by human societies. For our context of study, we show that it is possible to model human dependence on local natural resources with few demographic variables (population density, intensity of informal settlements, immigration rate and proportion of large size households) that can be produced at higher frequency and finer spatial resolution at minimum cost, by combining remote sensing with crowd-sourced information. The methodology is particularly suited for monitoring programs in data-scarce areas of the Global South, as a means to include estimates of ES use in addition to well-developed methods to understand ES supply, and can be generalized to meet the needs of different sustainability related issues. Fully understanding the spatial and temporal dynamics of human dependence on local natural resources is fundamental in order to target existing policies, or create new ones that can enable the maintenance of adequate local ecosystem health, even more so in rapidly urbanising contexts.

Acknowledgments

SB and FV are supported by the Basque Government through the BERC 2018-2021 program and by Spanish Ministry of Economy and Competitiveness MINECO through BC3 María de Maeztu excellence accreditation MDM-2017-0714. OS was supported by Sida funded GRAID project at Stockholm Resilience Centre. NS, RB, IK and POF were supported by the Council for Scientific and Industrial Research and SwedBio at the Stockholm Resilience Centre funded by Sida.

ORCID iDs

Stefano Balbi  <https://orcid.org/0000-0001-8190-5968>

Odirilwe Selomane  <https://orcid.org/0000-0002-6892-4221>

References

- Al-assaf A, Nawash O and Omari M 2014 Identifying forest ecosystem services through socio-ecological bundles: a case study from northern Jordan *Int. J. Sustain. Dev. World Ecol.* **21** 314–21
- Anderson K 1987 On why agriculture declines with economic growth *Agric. Econ.* **1** 195–207
- Anderson MJ 2006 Distance-based tests for homogeneity of multivariate dispersions *Biometrics* **62** 245–53
- Bertram C and Rehdanz K 2015 Preferences for cultural urban ecosystem services: comparing attitudes, perception, and use *Ecosyst. Serv.* **12** 187–99

- Bhorat H, Oosthuizen M and van der Westhuizen C 2012 Estimating a poverty line: an application to free basic municipal services in South Africa *Dev. South. Afr.* **29** 77–96
- Brouwer R, Akter S, Brander L and Haque E 2009 Economic valuation of flood risk exposure and reduction in a severely flood prone developing country *Environ. Dev. Econ.* **14** 397–417
- Brueckner J K and Lall S V 2015 Chapter 21—Cities in developing countries: fueled by rural-urban migration, lacking in tenure security, and short of affordable housing *Handbook Reg. Urban Econ.* **5** 1399–455
- Busgeeth M K, Brits A and Whisen J 2008 Potential application of remote sensing in monitoring informal settlements in developing countries where complimentary data does not exist *Planning Africa Conf. (Sandton Convention Centre, Johannesburg, South Africa, 14–16 April 2008)* (<https://researchspace.csir.co.za/dspace/handle/10204/3102>)
- Chakraborty A, Wilson B, Sarraf S and Jana A 2015 Open data for informal settlements: toward a user's guide for urban managers and planners *J. Urban Manage.* **4** 74–91
- Chandy L 2013 Counting the poor: methods, problems and solutions behind the \$1.25 a day global poverty estimates *Investments to End Poverty Working Paper Development Initiatives, Bristol* (<http://devinit.org/post/discussion-paper-counting-poor/>)
- Collinson M A, Tollman S M and Kahn K 2007 Migration, settlement change and health in post-apartheid South Africa: triangulating health and demographic surveillance with national census data *Scand. J. Publ. Health* **35** 77–84
- Cord A F, Brauman K A, Chaplin-Kramer R, Huth A, Ziv G and Seppelt R 2017 Priorities to advance monitoring of ecosystem services using Earth observation *Trends Ecol. Evol.* **32** 416–28
- Cord A F, Seppelt R and Turner W 2015 Sustainable development goals: monitor ecosystem services from space *Nature* **525** 33
- Cumming G S, Buerkert A, Hoffmann E M, Schlecht E, von Cramon-Taubadel S and Tschernitz T 2014 Implications of agricultural transitions and urbanization for ecosystem services *Nature* **515** 50–7
- Daunton M J 1995 *Progress and poverty: an economic and social history of Britain, 1700–1850* (Oxford: Oxford University Press) (<https://ideas.repec.org/b/oxp/obooks/9780198222811.html>)
- Daw T, Brown K, Rosendo S and Pomeroy R 2011 Applying the ecosystem services concept to poverty alleviation: the need to disaggregate human well-being *Environ. Conservation* **38** 370–9
- De Araujo Barbosa C C, Atkinson P M and Dearing J A 2015 Remote sensing of ecosystem services: a systematic review *Ecol. Indicators* **52** 430–43
- Djoudi H, Locatelli B, Vaast C, Asher K, Brockhaus M and Basnett Sijapati B 2016 Beyond dichotomies: gender and intersecting inequalities in climate change studies *Ambio* **45** 248–62
- Foster V and Briceno-Garmendia C 2010 *Africa's Infrastructure: A Time for Transformation: A Time for Transformation* (Washington, DC: The World Bank) (<https://openknowledge.worldbank.org/handle/10986/2692>)
- Grimm N B, Faeth S H, Golubiewski N E, Redman C L, Wu J, Bai X and Briggs J M 2018 Global change and the ecology of cities *Sci. J.* **319** 756–60
- Güneralp B, Lwasa S, Masundire H, Parnell S and Seto K C 2018 Urbanization in Africa: challenges and opportunities for conservation *Environ. Res. Lett.* **13** 015002
- Haider L J, Boonstra W J, Peterson G D and Schlüter M 2018 Traps and sustainable development in rural areas: a review *World Dev.* **101** 311–21
- Hamann M, Biggs R and Reyers B 2015 Mapping social-ecological systems: identifying 'green-loop' and 'red-loop' dynamics based on characteristic bundles of ecosystem service use *Glob. Environ. Change* **34** 218–26
- Hunt D V L *et al* 2012 Scenario archetypes: converging rather than diverging themes *Sustainability* **4** 740–72
- Karp D S *et al* 2015 National indicators for observing ecosystem service change *Glob. Environ. Change* **35** 12–21
- La Rosa D, Spyra M and Inostroza L 2016 *Indicators of Cultural Ecosystem Services for Urban Planning: A Review* (Amsterdam: Elsevier) pp 74–89
- Leach M, Mearns R and Scoones I 1999 Environmental entitlements: dynamics and institutions in community based natural resource management *World Dev.* **21** 225–47
- Liu J *et al* 2013 Framing sustainability in a telecoupled world *Ecol. Soc.* **18** art26
- Masterson V, Tengö M and Spierenburg M 2017 Competing place meanings in complex landscapes: a social-ecological approach to unpacking community conservation outcomes on the wild coast, South Africa *Soc. Nat. Resour.* **30** 1442–57
- McFarlane C 2012 Rethinking informality: politics, crisis, and the city *Plan. Theor. Pract.* **13** 89–108
- McHale M R, Bunn D N, Pickett S T and Twine W 2013 Urban ecology in a developing world: why advanced socioecological theory needs Africa *Frontiers Ecol. Environ.* **11** 556–64
- Mellor J W (ed) 1995 *Agriculture on the Road to Industrialization* (Baltimore, MD: The John Hopkins University Press)
- Millennium Ecosystem Assessment 2005 *Ecosystems and Human Well-Being: Synthesis* (Washington, DC: Island Press)
- Muller M 2008 *Free Basic Water—A Sustainable Instrument for a Sustainable Future in South Africa* (London: SAGE Publications Sage UK) pp 67–87
- Nel J L, Le Maitre D C, Roux D J, Colvin C, Smith J S, Smith-Adao L B, Maherry A and Sitas N 2017 Strategic water source areas for urban water security: making the connection between protecting ecosystems and benefiting from their services *Ecosyst. Serv.* **28** 251–9
- Newland K 2009 Circular migration and human development *UNDP Research Paper 2009/46* (New York: United Nations Development Programme) (http://hdr.undp.org/sites/default/files/hdrp_2009_42.pdf)
- Palomo I, Felipe-Lucia M R, Bennett E, Martín-López B and Pascual U 2016 Disentangling the pathways and effects of ecosystem service co-production *Adv. Ecol. Res.* **54** 245–83
- Pfeffermann D 2002 Small area estimation—new developments and directions *Int. Stat. Rev.* **70** 125–43
- Randall S and Coast E 2015 Poverty in African Households: the Limits of Survey and Census representations *J. Dev. Stud.* **51** 162–77
- Rao J N K and Yu M 1994 Small-area estimation by combining time-series and cross-sectional data *Can. J. Stat.* **22** 511–28
- Rao N, Lawson E T, Raditloaneng W N, Solomon D and Angula M N 2017 Gendered vulnerabilities to climate change: insights from the semi-arid regions of Africa and Asia *Clim. Dev.* **11** 14–26
- Raudsepp-Hearne C, Peterson G D and Bennett E M 2010 Ecosystem service bundles for analyzing tradeoffs in diverse landscapes *Proc. Natl Acad. Sci. USA* **107** 5242–7
- Reyers B, Biggs R, Cumming G S, Elmqvist T, Hejnowicz A P and Polasky S 2013 Getting the measure of ecosystem services: a social-ecological approach *Frontiers Ecol. Environ.* **11** 268–73
- Roy A 2009 Why India cannot plan its cities: informality, insurgence and the idiom of urbanization *Plan. Theor.* **8** 76–87
- Sanderson E W, Walston J and Robinson J G 2018 From bottleneck to breakthrough: urbanization and the future of biodiversity conservation *BioSci.* **68** 412–26
- Selomane O, Reyers B, Biggs R, Tallis H and Polasky S 2015 Towards integrated social-ecological sustainability indicators: exploring the contribution and gaps in existing global data *Ecol. Econ.* **118** 140–6
- Serajuddin U, Uematsu H, Wieser C, Yoshida N and Dabalen A 2015 Data deprivation—another deprivation to end *Policy Research Working Paper WPS 7252* (Washington, DC: The World Bank) (<http://documents.worldbank.org/curated/en/700611468172787967/Data-deprivation-another-deprivation-to-end>)
- Shackleton C, Shackleton S, Gambiza J, Nel E, Rowntree K and Urquhart P 2008 Links between ecosystem services and poverty alleviation: situation analysis for arid and semi-arid

- lands in southern Africa 2008 *Ecosystem Services and Poverty Reduction Research Programme: DFID, NERC, ESRC* (submitted)
- Soubbotina T P and Sheram K 2000 *Beyond Economic Growth: Meeting the Challenges of Global Development* (in English) (Washington, DC: World Bank) (<http://documents.worldbank.org/curated/en/408121468149372828/Beyond-economic-growth-meeting-the-challenges-of-global-development>)
- StatsSA 2011 *Census 2011 metadata* Statistics South Africa, Pretoria
- StatsSA 2012 *Poverty trends in South Africa: an examination of absolute poverty between 2006 and 2011* Statistics South Africa, Pretoria
- Syrbe R U and Walz U 2012 Spatial indicators for the assessment of ecosystem services: providing, benefiting and connecting areas and landscape metrics *Ecol. Indicators* **21** 80–8
- Tadesse G, Zavaleta E, Shennan C and FitzSimmons M 2014 Local ecosystem service use and assessment vary with socio-ecological conditions: a case of native coffee-forests in southwestern Ethiopia *Human Ecol.* **42** 873–83
- Tallis H *et al* 2012 A global system for monitoring ecosystem service change *BioSci.* **62** 977–86
- UN 2015 *Transforming our World: the 2030 Agenda for Sustainable Development* (New York: United Nations) (<https://sustainabledevelopment.un.org/post2015/transformingourworld>)
- Unterhalter E 2012 Poverty, education, gender and the millennium development goals: reflections on boundaries and intersectionality *Theory Res. Educ.* **10** 253–74
- Willcock S *et al* 2018 Machine learning for ecosystem services *Ecosyst. Serv.* **33** 165–74
- Witten I H, Frank E, Hall M A and Pal C J 2018 *Data Mining: Practical Machine Learning Tools and Techniques* 4 ed (San Mateo, CA: Morgan Kaufmann Publishers)
- Xie M, Jean N, Burke M, Lobell D B and Ermon S 2015 Transfer learning from deep features for remote sensing and poverty mapping *Proc. 30th AAAI Conf. on Artificial Intelligence* (arXiv:1510.00098)
- Yang H, Harrison R, Yi Z-F, Goodale E, Zhao M-X and Xu J-C 2015 Changing perceptions of forest value and attitudes toward management of a recently established nature reserve: a case study in Southwest China *Forests* **6** 3136–64
- Yang W, Dietz T, Kramer D B, Chen X and Liu J 2013 Going beyond the millennium ecosystem assessment: an index system of human well-being *PLoS ONE* **8** e64582